

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 1 026 263 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

**09.08.2000 Bulletin 2000/32**

(51) Int. Cl.<sup>7</sup>: **C12Q 1/70**

(21) Application number: **00300792.9**

(22) Date of filing: **01.02.2000**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **02.02.1999 US 118417 P**

(71) Applicant:

**Ortho-Clinical Diagnostics, Inc.  
Rochester, NY 14626-5101 (US)**

(72) Inventors:

• **Patterson, David R.  
San Diego, California 92128 (US)**

• **Puskas, John A.**

**Rochester, New York 14613 (US)**

• **Song, Keming**

**Ballwin, Missouri 63021 (US)**

• **Linnen, Jeffrey M.**

**San Diego, California 92129 (US)**

(74) Representative:

**Fisher, Adrian John**

**CARPMAELS & RANSFORD**

**43 Bloomsbury Square**

**London WC1A 2RA (GB)**

Remarks:

The applicant has subsequently filed a sequence listing and declared, that it includes no new matter.

(54) **Oligonucleotide reverse transcription primers for efficient detection of HIV-1 and HIV-2 and methods of use thereof**

(57) Disclosed herein are methods and kits for the detection of human immunodeficiency virus in biological samples from human subjects. Oligonucleotide reverse transcription primers for use in such methods and kits for detection of human immunodeficiency virus are also described.

**EP 1 026 263 A2**

## Description

### Field of the Invention

- 5 [0001] The present invention pertains to improved methods for detecting nucleic acid sequences in biological samples, particularly sequences derived from infectious microorganisms.

### Background of the Invention

- 10 [0002] Millions of individuals world-wide are infected with Human Immunodeficiency Virus (HIV). Consequently, HIV infection represents a serious public health concern. Spread of HIV infection via contaminated blood products means that there is a need for screening methods that can detect small amounts of HIV RNA in patient samples. Furthermore, the increasing availability of ameliorative treatments for HIV infection means that early detection of infection in a patient is vital in order to initiate appropriate therapeutic interventions.
- 15 [0003] Thus, there is a need in the art for highly sensitive detection methods for HIV that can be used in diagnosis and screening.

### Summary of the Invention

- 20 [0004] The present invention provides a method for reverse transcribing Human Immunodeficiency Virus (HIV) RNA in a biological sample, where the method comprises:

(a) contacting RNA derived from said sample with an oligonucleotide under conditions in which said oligonucleotide primes synthesis of DNA complementary to at least a portion of said RNA;

- 25 wherein said oligonucleotide is selected from the group consisting of

- (i) 5'-CTTGTATTACTACTG-3' (SEQ ID NO 1),  
 (ii) 5'-CCCTGTGGCGCC-3' (SEQ ID NO 2),  
 (iii) 5'-GCGACTAGGAGAGA-3' (SEQ ID NO 3),  
 30 (iv) 5'-CCCAGACGGTCAGT-3' (SEQ ID NO 4), or  
 (v) any combination of any of the foregoing.

- [0005] In another aspect, the invention provides a method for detecting the presence of Human Immunodeficiency Virus (HIV) RNA in a biological sample, where the method comprises:

- 35 (a) performing a reverse transcription reaction using as a template RNA derived from the sample and using, as a primer, an oligonucleotide complementary to a nucleotide sequence contained within the RNA to produce HIV-specific reverse transcription products,

where the primer is selected from the group consisting of:

- 40 (i) 5'-CTTGTATTACTACTG-3' (SEQ ID NO 1),  
 (ii) 5'-CCCTGTGGCGCC-3' (SEQ ID NO 2),  
 (iii) 5'-GCGACTAGGAGAGA-3' (SEQ ID NO 3),  
 (iv) 5'-CCCAGACGGTCAGT-3' (SEQ ID NO 4), or  
 45 (v) any combination of any of the foregoing;

(b) amplifying products of the reverse transcription reaction to produce amplification products; and

(c) detecting the amplification products;

where detection of the amplification products indicates the presence of HIV RNA in the sample.

- 50 [0006] Amplification may be carried out by any method, preferably polymerase chain reaction (PCR). The use of HIV-specific reverse transcription primers according to the invention provides a sensitive method for detecting HIV-1 and/or HIV-2 in a sample, preferably plasma.

- [0007] In yet another aspect, the invention provides kits for the detection of HIV-1, HIV-2, or a combination thereof  
 55 in a biological sample, where the kit comprises a reverse transcription primer selected from the group consisting of:

- (a) 5'-CTTGTATTACTACTG-3' (SEQ ID NO 1),  
 (b) 5'-CCCTGTGGCGCC-3' (SEQ ID NO 2),

- (c) 5'-GCGACTAGGAGAGA-3' (SEQ ID NO 3),  
 (d) 5'-CCCAGACGGTCAGT-3' (SEQ ID NO 4), or  
 (e) any combination of any of the foregoing. The kits may additionally comprise reagents and instructions for reverse transcription, amplification, and product detection.

5

### **Brief Description of the Drawings**

[0008]

- 10 Figure 1 is a photographic illustration of a 4% agarose gel stained with ethidium bromide showing HIV-1-specific amplification products obtained using as a reverse transcription primer (a) random hexamer primers (lanes 2-10); or (b) a mixture of random hexamer primers and LTR8RT (lanes 12-20). Lane 1 contains markers, and lanes 22, 24, and 26 are control samples.
- 15 Figure 2 is a photographic illustration of a 4% agarose gel stained with ethidium bromide showing HIV-1-specific amplification products obtained using as a reverse transcription primer (a) a mixture of random hexamer primers and POL3RT (lanes 2-10) or (b) a mixture of LTR8RT and POL3RT (lanes 12-20). Lane 1 contains markers, and lanes 22, 24, and 26 are control samples.
- 20 Figure 3 is a photographic illustration of a 4% agarose gel stained with ethidium bromide showing HIV-1-specific amplification products obtained using (a) random hexamer primers (lanes 2-13) or (b) random hexamer and a mixture of POL3RT, LTR8RT, 2LTRRT and 2EnvRT (lanes 15-26). Lane 1 contains markers.
- Figure 4 is a photographic illustration of a 4% agarose gel stained with ethidium bromide showing HIV-1-specific amplification products obtained using (a) random hexamer primers (lanes 2-15) or (b) random hexamer and a mixture of POL3RT, LTR8RT, 2LTRRT and 2EnvRT (lanes 16-29). Lanes 1 and 30 contain markers.

### **Detailed Description of the Invention**

- 25 [0009] The present inventors have discovered that detection of Human Immunodeficiency Virus (HIV) RNA in biological samples is more efficient when oligonucleotides having sequences complementary to certain sequences present in HIV RNA are used as primers for reverse transcription. Preferably, the sequences of the primers correspond to sequences near the 3' end of HIV RNA.

- 30 [0010] Many techniques in molecular biology, microbiology, recombinant DNA, and protein biochemistry are used in practicing the present invention, such as those explained in, for example, *Current Protocols in Molecular Biology*, Volumes I, II, and III, 1997 (F.M. Ausubel ed.); Sambrook *et al.*, 1989, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; *DNA Cloning: A Practical Approach*, Volumes I and II, 1985 (D.N. Glover ed.); *Oligonucleotide Synthesis*, 1984, (M.L. Gait ed.); *Transcription and Translation*, 1984 (Hames and Higgins eds.); *A Practical Guide to Molecular Cloning*; the series, *Methods in Enzymology* (Academic Press, Inc.); and *Protein Purification: Principles and Practice*, Second Edition (Springer-Verlag, N.Y.).

- 35 [0011] "Nucleic acid" or "polynucleotide" as used herein refers to purine- and pyrimidine-containing polymers of any length, either polyribonucleotides or polydeoxyribonucleotides or mixed polyribo-polydeoxyribo nucleotides. This includes single- and double-stranded molecules, such as, for example, DNA-DNA, DNA-RNA and RNA-RNA hybrids, as well as "protein nucleic acids" (PNA) formed by conjugating bases to an amino acid backbone. This also includes nucleic acids containing modified bases.

- 40 [0012] A "complement" of a nucleic acid sequence as used herein refers to the antisense sequence that participates in Watson-Crick base-pairing with the original sequence.

- 45 [0013] A "primer" as used herein is an oligonucleotide between about 5 and about 50 nucleotides in length, preferably between about 6 and about 25 nucleotides in length and most preferably between about 6 and about 18 nucleotides in length, that forms a duplex with a single-stranded nucleic acid sequence of interest and allows polymerization of a complementary strand using, e.g., reverse transcriptase or DNA polymerase.

- 50 [0014] An "isolated" nucleic acid or polypeptide as used herein refers to a component that is removed from its original environment (for example, its natural environment if it is naturally occurring or a reaction mixture if it is synthetic). An isolated nucleic acid or polypeptide typically contains less than about 50%, preferably less than about 75%, and most preferably less than about 90%, of the components with which it was originally associated.

- [0015] A nucleic acid sequence that is "derived from" a designated sequence refers to a sequence that corresponds to a region of the designated sequence. This encompasses sequences that are homologous or complementary to the sequence.

- 55 [0016] An internal positive control (IPC) target nucleic acid refers to a synthetic nucleic acid sequence cloned into a plasmid vector which is subsequently linearized, typically by the action of a restriction endonuclease. An IPC will typically have multiple primer binding sequences surrounding a generic probe-binding region, and acts as a generic control

against false negative results in nucleic acid amplification reactions.

**[0017]** The sequence of a preferred internal positive control target DNA is:

5'-CGCCAGCGTGGACCATCAAGTAGTAATGAACGCACGGACGAGGACATCA TAGAGATTACACCTTTATCCACAGT-  
TCTCGGTCTAACGCAGCAGTCAGTG TATCAGCACCAGCATCCGTAGTGAGTCTTCAGTGTCTGCTCCAGGATCGT  
G-3' (SEQ ID NO 5).

**[0018]** As used herein, conditions appropriate for reverse transcription, i.e., conditions in which an oligonucleotide will prime cDNA synthesis, encompass incubation of RNA and primer oligonucleotides with a reverse transcriptase enzyme and nucleotides at a temperature and for a time that results in synthesis of cDNA.

**[0019]** Nucleic acids comprising any of the sequences disclosed herein or subsequences thereof can be prepared by conventional methods. For example, DNA can be chemically synthesized using, e.g., the phosphoramidite solid support method of Matteucci *et al.*, 1981, *J. Am. Chem. Soc.* **103**:3185, the method of Yoo *et al.*, 1989, *J. Biol. Chem.* **264**:17078, or other well known methods. The nucleic acids may also be modified by many means known in the art. Non-limiting examples of such modifications include methylation, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, and internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.) or charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.). Nucleic acids may contain one or more additional covalently linked moieties, such as, for example, proteins (e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), intercalators (e.g., acridine, psoralen, etc.), chelators (e.g., metals, radioactive metals, iron, oxidative metals, etc.), and alkylators. PNAs are also encompassed by the term "nucleic acid". The nucleic acid may be derivatized by formation of a methyl or ethyl phosphotriester or an alkyl phosphoramidate linkage. Furthermore, the nucleic acid sequences of the present invention may also be modified with a label capable of providing a detectable signal, either directly or indirectly. Exemplary labels include radioisotopes, fluorescent molecules, biotin, and the like.

**[0020]** Amplification as used herein refers to an iterative process by which a nucleic acid is copied. Suitable methods for amplification include without limitation the polymerase chain reaction, the ligase chain reaction, and transcription-mediated amplification.

**[0021]** Human Immunodeficiency Virus (HIV) as used herein refers to species in the genus of Retroviridae, including HIV-1, HIV-2, and SIV, and variant strains thereof. Isolates of HIV that may be detected by the present invention include, but are not limited to, HIV-1 and HIV-2.

**[0022]** The present invention provides methods for reverse transcribing HIV RNA from biological samples, which methods are useful for detection of HIV in biological samples. Detection of HIV-specific amplification products indicates the presence of HIV RNA in the sample.

**[0023]** According to the invention, a biological sample is obtained from a patient by any conventional means. Suitable biological samples include, without limitation, blood, serum, plasma, urine, breast milk, tissue samples, and cerebrospinal fluid. Preferably, plasma is used as the source of HIV RNA.

**[0024]** The biological sample is treated in any manner that provides access of the reverse transcription reagents to RNA, specifically HIV RNA, contained within the sample. RNA "derived from" a biological sample is any RNA which was originally present in the sample and to which access has been gained by treating the sample. Preferably, RNA is extracted from the sample using any method well known in the art, such as, e.g., methods employing guanidinium thiocyanate, or using commercially available reagents and methods such as, e.g., PureScript<sup>®</sup> from Gentra Systems, Inc. (Minneapolis MN). Any extraction procedure may be used that results in separation from the RNA of RNases, other proteins, and/or any other components that might interfere with reverse transcription.

**[0025]** The RNA extracted from the sample is then contacted with oligonucleotide primers under conditions where the oligonucleotides prime the synthesis of DNA complementary to at least a portion of the extracted RNA. The sequences of the oligonucleotide primers are derived from the sequence of HIV. The primers correspond to regions of HIV RNA that may be downstream, i.e., 3', to regions whose detection is desired. These regions may include, e.g., the long terminal repeat (LTR) region, the region encoding the viral reverse transcriptase (Pol), Gag protein, Tat protein, envelope glycoprotein, Vif, Vpr, and Vpu proteins, and the Rev region, which encodes a transcription factor response element. Preferably, the primers correspond to sequences near the 3' end of the HIV genome. The primer sequences may be used to specifically identify particular isolates of HIV (e.g., isolates of HIV-1 and HIV-2). A primer may identify a particular isolate by hybridizing to RNA derived from that isolate under conditions in which it does not hybridize to RNA from a different isolate, i.e., the primer itself may comprise a sequence that differs between isolates. Alternatively, the primer sequence may be used to prime synthesis of a fragment of HIV RNA that differs between isolates, i.e., the sequence that differs between the isolates may be downstream of the primer sequence.

**[0026]** Reverse transcription primers useful in practicing the present invention are selected based on theoretical considerations of sequence conservation, intra- and inter-molecular interactions, and the predicted secondary structures of the amplicon and surrounding sequence. Furthermore, the primers and assay system are designed to allow the co-amplification (and co-detection) of multiple regions of the HIV genome, multiple viral species, and an internal positive control (IPC) RNA (or DNA).

[0027] Non-limiting examples of reverse transcription primers according to the invention are shown in Table 1.

TABLE 1

SOURCE	DESIGNATION	SEQUENCE	SEQ ID NO.
HIV-1	POL3RT	5'-CTTGTTACTACTG-3'	1
HIV-1	LTR8RT	5'-CCCTGTGGCGCC-3'	2
HIV-2	2LTR1RT	5'-GCGACTAGGAGAGA-3'	3
HIV-2	2Env2RT	5'-CCCAGACGGTCAGT-3'	4

[0028] Reverse transcription is performed using one or more of the above primers. Random primers, such as, e.g., random hexamer reverse transcription primers (N6, Pharmacia Biotech, Piscataway, NJ) may also be added. Reverse transcription is carried out using conventional procedures, such as are described in *Current Protocols in Molecular Biology*, Volumes I, II, and III, 1997 (F.M. Ausubel ed.); in U.S. Patent 5,322,770; in Young, et al., *J. Clin. Microbiol.* 31(4):882 (1993); Myers et al., *Biochemistry* 30(3):7661 (1991); or as described in copending patent application No. \_\_\_\_\_ (ref CDS-212), claiming priority of USSN 60/118,520.

[0029] Following the reverse transcription reaction, the cDNA product or products can be isolated and recovered by conventional methods. Preferably, the cDNA product or products are amplified. Any method for amplification may be used, including, without limitation, polymerase chain reaction (PCR), ligase chain reaction, strand displacement amplification, transcript mediated amplification, and nucleic acid single base amplification. Preferably, PCR is used. Typically, a reaction mixture containing all of the necessary components for PCR (including HIV-specific amplification primers) is added directly to the reverse transcription reaction mixture. Amplification is then carried out using conditions specified by the primer pairs that are used. Suitable amplification primer pairs are disclosed, e.g. in copending patent application No. \_\_\_\_\_ (ref CDS-210), claiming priority of USSN 60/118,498, and in the Examples below.

[0030] Following amplification, the amplification products may be detected using any method known in the art, including, without limitation, gel electrophoresis in agarose or acrylamide; capture of the amplification products on a solid support followed by colorimetric detection (see, e.g., Example 1 below); ECI detection; fluorescence, radioisotopic detection, and chemiluminescence. Reagents for such detection methods are commercially available from, e.g. Molecular Probes, Eugene, Oregon and Ortho Clinical Diagnostics, Rochester, NY.

[0031] The detection of HIV-specific amplification products indicates the presence of HIV RNA in the sample. When gel electrophoresis is used, HIV-specific amplification products are confirmed by their size, as predicted by the location in HIV RNA of the sequences corresponding to the amplification primers used in the reaction.

[0032] The present invention provides kits for detection of HIV RNA in biological samples, which comprise one or more of the reverse transcription primers shown in Table 1 above. The kits may also comprise reagents for reverse transcription, as well as additional reagents for detection of HIV cDNA by, e.g., PCR.

#### **Description of the Preferred Embodiments**

[0033] The following examples illustrate the present invention without limitation.

##### Methods:

##### 1. Sample preparation:

[0034] RNA was prepared from plasma samples using guanidinium thiocyanate or PureScript<sup>®</sup> RNA isolation reagents (Gentra Systems, Minneapolis MN). Modifications to the manufacturer's protocol for body fluids included use of 40µg glycogen, rather than 20µg, as a carrier to aid in the precipitation of viral RNA. Additionally, in most cases, after isopropyl alcohol precipitation of the RNA and washing the RNA pellet with ethanol, the RNA pellet was resuspended in the RT buffer mix, rather than in the RNA hydration solution provided by the manufacturer.

##### 2. Reverse Transcription:

[0035] The synthesis of cDNA from RNA was catalyzed by the addition of 100 U recombinant Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase (RT) (Gibco BRL, Gaithersburg, Maryland) in a 50µl solution of 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl<sub>2</sub>, 10 mM DTT, 0.4 mM of each dNTP (Pharmacia Biotech), 4µM random hex-

amers (Pharmacia Biotech, Piscataway, NJ) and/or specific reverse transcription primer, and 20 units RNasin (Promega, Madison, Wisconsin) in diethylpyrocarbonate (DEPC)-treated water. After incubation at 42°C for 30 min, the RT reaction was held at 100°C for 5 min to destroy RT activity. Each reaction was chilled for 1 min followed by micro-centrifugation at 16000 x g for 4 seconds.

### 3. PCR amplification:

**[0036]** PCR was carried out in a PE9600 thermocycler (Perkin-Elmer) in a 100 µl solution of 25 mM Tris-HCl, 3 mM MgCl<sub>2</sub>, 0.725 mM EDTA, 54 mM KCl, 3.72 mM NaCl, 40 µM DTT, 108 µg/mL gelatin (type IV), 9.5% glycerol, 0.02% Tween 20, 0.02% NP40, calf thymus DNA (2 µg), 1.2 mM of each dNTP, 0.4 µM of each primer, 10 copies linearized internal positive control (IPC) plasmid DNA, and 16 U of Taq polymerase. Monoclonal antibodies to Taq, TP1-12 and TP4-9, the preparation of which are disclosed in U.S. Patent 5,338,671, were added to the reaction at a 50:1 and 5:1 molar ratio, respectively, to provide a 55:1 molar ratio of antibody to Taq polymerase. After initial denaturation at 96°C for 3 min, 40 cycles of amplification were performed at 96°C for 5 sec and 68°C for 40 sec. At the conclusion of cycling, a post-heat step was performed for 5 min at 103°C to inactivate Taq polymerase. The amplification primers used are shown in Table 2 below.

TABLE 2

ID	Source	Sequence	SEQ ID NO.
JBLT R4	HIV-1- (s)	5'-CTG CTT AAG CCT CAA TAA AGC TTG CCT TGA-3'	6
JBLT R6	HIV-1(as)	5'-GGG TCT GAG GGA TCT CTA GTT ACC AGA GT-3'	7
JBLT R8	HIV-1(as)	5'-TGT TCG GGC GCC ACT GCT AGA GA-3'	8
2LTRe	HIV-2 (s)	5'-GGG AGG TTC TCT CCA GCA CTA GCA-3'	9
2LTR-R1	HIV-2 (as)	5'-GCG ACT AGG AGA GAT GGG AAC ACA CA-3'	10

### 4. Detection of PCR products:

**[0037]** PCR products were detected either by (i) gel electrophoresis, followed by ethidium bromide staining; or (ii) use of 5'-biotin-labeled primers (sense strand) during amplification. In this case, the amplification products were captured by hybridization to oligonucleotide probes covalently attached to latex particles, which were deposited on the surface of a flow through membrane (SureCell® tests, Ortho Clinical Diagnostics, Rochester, NY). The HIV-1 probes were: 5'-CAACAGACGGGCACACACTACT-3'(JBLTRpr) (SEQ ID NO 11) and 5'-GAACAGATGGGCACACACTGCT-3'(JBLTRpr4) (SEQ ID NO 12); and the HIV-2 probe was 5'-CCACGCTTGCTTGCTTAAAGACCTC-3'(2LTRpr1) (SEQ ID NO 13). The probe/product complex was reacted with streptavidin (SA)-horseradish peroxidase (HRP) conjugate, which catalyzes the oxidative conversion of a dye precursor to a dye (blue color). The blue color intensity was scored visually (0-10) by comparing color intensity to color standards. All visual color scores > 3 were considered to be positive results.

#### Example 1: Efficiency of Reverse Transcription Using HIV-Specific Primers or Random Primers

**[0038]** The following experiment was performed to compare the efficiency of reverse transcription of HIV RNA derived from human plasma samples using HIV-specific primers according to the invention or random hexamer reverse transcription primers (N6, Pharmacia Biotech).

**[0039]** Human plasma was diluted to contain 1000 copies of HIV RNA per 100 µl, to yield approximately 100 copies of HIV RNA per reaction. RNA was extracted from the plasma using guanidinium thiocyanate. The RNA pellet was dissolved in 26 µl of diethylpyrocarbonate-treated water.

**[0040]** The reverse transcription reaction contained: 13 µl RNA, 10 µl reverse transcription mix (which contained first-strand buffer, 0.1 M DTT, 20 U RNasin (Promega, Madison, WI), 0.4 mM of each dNTP, and 200 Units of Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase). An additional 2 µl of the following primer mixes (all at 50 µM) were added according to the condition being tested: (1) 2 µl N6 random primers; (2) 1 µl N6 random primers + 1 µl LTR8RT primer; (3) 1 µl N6 random primers + 1 µl POL3RT primer; (4) 1 µl LTR8RT + 1 µl POL3RT. The reverse transcription reaction was incubated at 42°C for 30 minutes; heated to 100°C for 5 minutes; and then chilled on ice for 1 minute. A 75 µl PCR master mix was then added to the cDNA-containing reaction mixture and PCR was performed under the following conditions: a 3 minute preheat at 96°C, followed by 5 cycles of melting at 96 °C followed by anneal-

ing and amplification at 62°C for 5 seconds, followed by 35 cycles of melting at 96°C and annealing and amplifying at 68°C for 40 seconds. The amplification products were then resolved in a 4% agarose gel and visualized using ethidium bromide.

[0041] *Results:* As illustrated in Figures 1 and 2, the use of HIV-specific reverse transcription primers, either alone or in conjunction with random hexamer primers, results in the detection of significantly more HIV-1-specific amplification products. Compare Figure 1, lanes 2-10 (random primer alone) with Figure 1, lanes 12-20 (LTR8RT + random primer); Figure 2, lanes 2-10 (POL3RT + random primer); and Figure 2, lanes 12-20 (LTR8RT + POL3RT). This result was also observed in when 100  $\mu$ M or 200  $\mu$ M random primers were used.

#### Example 2: Detection of HIV RNA in Patient Samples

[0042] The following study was performed to compare the detection of HIV RNA in patient samples using either random hexamer reverse transcription primers or random primers in conjunction with HIV -specific reverse transcription primers.

[0043] HIV-positive plasma samples were collected from patients having CD4 T-cell counts greater than 500, indicating that they were asymptomatic and had a relatively low viral load.

[0044] RNA was extracted from the plasma samples as described in Example 1 above. 13  $\mu$ l of the RNA solution were diluted in 15  $\mu$ l water. Each sample was split into two 12- $\mu$ l aliquots for reverse transcription. Two reverse transcription reaction mixes were prepared with as described in Example 1 above. Each mix contained either 2  $\mu$ l of 100  $\mu$ M random primer + 1  $\mu$ l water or 2  $\mu$ l of 100  $\mu$ M random primer + 1  $\mu$ l of a 50  $\mu$ M HIV -specific primer mix containing equal amounts of the following primers: (1) POL3RT; (2) LTR8RT; (3) 2LTRRT; and (4) 2EnvRT. The reverse transcription and amplification reactions were performed as described in Example 1 above.

[0045] HIV-specific amplification products were detected by gel electrophoresis on 4% agarose gels stained with Ethidium Bromide and also by the SureCell<sup>®</sup> colorimetric method described above.

[0046] *Results:* Figures 3 and 4 illustrate the amplification products detected by gel electrophoresis. Detection of HIV-specific amplification products by the colorimetric method is indicated in relative values in Table 3. IPC indicated internal positive control primers.

TABLE 3

	RANDOM PRIMERS ONLY			N6 + RT PRIMERS		
	LTR	POL	IPC	LTR	POL	IPC
Sample	3/4	3/4	1P	3/4	3/4	1P
1	0	0	8	0	0	8
2	8	7	8	8	7	7.5
3	5	6	8	7	7	7.5
4	5	5	8	5	5	8
5	7	5	8	7.5	7.5	8
6	na	na	na	na	na	na
7	6	7	8	5	7.5	8
8	8	8	8	8	8	8
9	2	2	8	3	0	8
10	5	7	8	7	8	8
11	6	7	8	6	6	8
12	8	8.5	8	7.5	9	7.5
13	8	7.5	8	8	7.5	8
14	0	0	8	0	0	9
15	0	0	8	0	0	8
16	2	0	8	7	8	9



TABLE 3 (continued)

	RANDOM PRIMERS ONLY			N6 + RT PRIMERS		
	LTR	POL	IPC	LTR	POL	IPC
Sample	3/4	3/4	1P	3/4	3/4	1P
17	7.5	7.5	8	7.5	7.5	8
18	6	7	8	6	7	8
19	7.5	7.5	7.5	7.5	8	8
20	2	1	8	5	5	8
21	6.5	6.5	8	7	7.5	7
22	1	1	8	3	5	8
23	3	1	8	2	2	8
24	7	9	8	7	9	7
25	na	na	na	na	na	na
26	9	9	7	9	9	6.5
27	3	4	7	5	7	8.5
28	2	2	8	2	2	8.5
Neg	0	0	7	0	0	7
Pos	5	5	7	5	5	7

[0047] In samples 3 and 10, the addition of HIV-specific RT primers to the random hexamer primers resulted in a greater degree of amplification than using random primers alone. The amount of product detected by the colorimetric method was also greater in these samples when HIV-specific RT primers were used.

[0048] Samples 16, 20, and 22 were not detected using only random primers in the reverse transcription reaction, either by gel electrophoresis or by colorimetry. These samples, however, were positive when HIV-specific RT primers were used in addition to random primers.

[0049] These results indicate that the methods and compositions of the present invention can reduce the incidence of false negative results in screening of patients or blood supply for HIV.

[0050] All patents, applications, articles, publications, and test methods mentioned above are hereby incorporated by reference in their entirety.

[0051] Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed description. Such obvious variations are within the full intended scope of the appended claims.

# EP 1 026 263 A2

## SEQUENCE LISTING

5 <110> ORTHO-CLINICAL DIAGNOSTICS, INC.  
 <120> OLIGONUCLEOTIDE REVERSE TRANSCRIPTION PRIMERS FOR EFFICIENT  
 DETECTION OF HIV-1 AND HIV-2 AND METHODS OF USE THEREOF

<130> P023846EP: AJF

10 <140> 00300792.9  
 <141> 01.02.00

<150> US 60/118417  
 <151> 02.02.99

15 <160> 13

<170> SeqWin99

<210> 1  
 <211> 15  
 20 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PRIMER

25 <400> 1  
 cttgtattac tactg 15

<210> 2  
 <211> 12  
 <212> DNA  
 30 <213> Artificial Sequence

<220>  
 <223> PRIMER

35 <400> 2  
 ccctgtggcg cc 12

<210> 3  
 <211> 14  
 <212> DNA  
 40 <213> Artificial Sequence

<220>  
 <223> PRIMER

45 <400> 3  
 gcgactagga gaga 14

<210> 4  
 <211> 14  
 <212> DNA  
 <213> Artificial Sequence

50 <220>  
 <223> PRIMER

<400> 4

55

# EP 1 026 263 A2

cccagacggt cagt 14

5 <210> 5  
 <211> 150  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PRIMER

10 <400> 5  
 cgccagcgtg gaccatcaag tagtaatgaa cgcacggacg aggacatcat agagattaca 60  
 cctttatcca cagttctcgg tctaacgcag cagtcagtg atcagcacca gcatccgtag 120  
 tgagtcttca gtgtctgctc caggatcgtg 150

15 <210> 6  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PRIMER

20 <400> 6  
 ctgcttaagc ctcaataaag cttgccttga 30

25 <210> 7  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PRIMER

30 <400> 7  
 ggggtctgagg gatctctagt taccaga 27

35 <210> 8  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PRIMER

40 <400> 8  
 tgttcgggcg ccactgctag aga 23

45 <210> 9  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PRIMER

50 <400> 9  
 gggagggttct ctccagcact agca 24

<210> 10  
 <211> 26

55

<212> DNA  
 <213> Artificial Sequence  
 5 <220>  
 <223> PRIMER  
 <400> 10  
 gcgactagga gagatgggaa cacaca 26  
 10 <210> 11  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence  
 15 <220>  
 <223> CAPTURE PROBE  
 <400> 11  
 20 caacagacgg gcacacacta ct 22  
 <210> 12  
 <211> 22  
 <212> DNA  
 25 <213> Artificial Sequence  
 <220>  
 <223> CAPTURE PROBE  
 <400> 12  
 30 gaacagatgg gcacacactg ct 22  
 <210> 13  
 <211> 25  
 <212> DNA  
 35 <213> Artificial Sequence  
 <220>  
 <223> CAPTURE PROBE  
 <400> 13  
 40 ccacgcttgc ttgcttaaag acctc 25

45 **Claims**

1. A method for reverse transcribing Human Immunodeficiency Virus (HIV) RNA in a biological sample, said method comprising:

50 (a) contacting RNA derived from said sample with an oligonucleotide under conditions in which said oligonucleotide primes synthesis of DNA complementary to at least a portion of said RNA;  
 wherein said oligonucleotide is selected from the group consisting of:  
 55 (i) 5'-CTTGTATTACTACTG-3' (SEQ ID NO 1),  
 (ii) 5'-CCCTGTGGCGCC-3' (SEQ ID NO 2),  
 (iii) 5'-GCGACTAGGAGAGA-3' (SEQ ID NO 3),  
 (iv) 5'-CCCAGACGGTCAGT-3' (SEQ ID NO 4), and

(v) any combination of any of the foregoing.

2. A method as defined in claim 1, wherein said sample is selected from the group consisting of blood, serum, plasma, urine, saliva, and cerebrospinal fluid.

3. A method as defined in claim 1, further comprising (b) recovering said cDNA.

4. A method for detecting the presence of Human Immunodeficiency Virus (HIV) RNA in a biological sample, said method comprising:

(a) performing a reverse transcription reaction, using RNA derived from said sample as a template and using an oligonucleotide complementary to a nucleotide sequence contained within said RNA as a reverse transcription primer to produce HIV-specific reverse transcription products, wherein said reverse transcription primer is selected from the group consisting of:

- (i) 5'-CTTGTATTACTACTG-3' (SEQ ID NO 1),
- (ii) 5'-CCCTGTGGCGCC-3' (SEQ ID NO 2),
- (iii) 5'-GCGACTAGGAGAGA-3' (SEQ ID NO 3),
- (iv) 5'-CCCAGACGGTCAGT-3' (SEQ ID NO 4), and
- (v) any combination of any of the foregoing;

(b) amplifying said reverse transcription products to produce amplification products; and

(c) detecting said amplification products;

wherein detection of said amplification products of said amplification indicates the presence of HIV RNA in said sample.

5. A method as defined in claim 4, wherein said sample is selected from the group consisting of blood, serum, plasma, urine, saliva, and cerebrospinal fluid.

6. A method as defined in claim 4, wherein said amplifying is performed by a method selected from the group consisting of polymerase chain reaction, ligase chain reaction, and strand displacement amplification.

7. A method as defined in claim 4, wherein said detecting is performed by a method selected from the group consisting of gel electrophoresis of amplification products, capture of amplification products on solid supports, and chemiluminescent detection of amplification products.

8. The oligonucleotide 5'-CTTGTATTACTACTG-3' (SEQ ID NO 1).

9. The oligonucleotide 5'-CCCTGTGGCGCC-3' (SEQ ID NO 2).

10. The oligonucleotide 5'-GCGACTAGGAGAGA-3' (SEQ ID NO 3).

11. The oligonucleotide 5'-CCCAGACGGTCAGT-3' (SEQ ID NO 4).

12. An HIV-specific reverse transcription primer comprising the oligonucleotide 5'-CTTGTATTACTACTG-3' (SEQ ID NO 1).

13. An HIV-specific reverse transcription primer comprising the oligonucleotide 5'-CCCTGTGGCGCC-3' (SEQ ID NO 2).

14. An HIV-specific reverse transcription primer comprising the oligonucleotide 5'-GCGACTAGGAGAGA-3' (SEQ ID NO 3).

15. An HIV-specific reverse transcription primer comprising the oligonucleotide 5'-CCCAGACGGTCAGT-3' (SEQ ID NO 4).

16. A kit for detection of HIV-1, HIV-2, or a combination thereof, in a biological sample, said kit comprising a reverse transcription primer selected from the group consisting of:

- (a) 5'-CTTGTATTACTACTG-3' (SEQ ID NO 1),  
(b) 5'-CCCTGTGGCGCC-3' (SEQ ID NO 2),  
(c) 5'-GCGACTAGGAGAGA-3' (SEQ ID NO 3),  
(d) 5'-CCCAGACGGTCAGT-3' (SEQ ID NO 4),  
(e) any combination of any of the foregoing.

5

17. A method as defined in claim 1, further comprising simultaneously contacting RNA derived from said sample with random hexamer oligonucleotides under conditions in which said random hexamer oligonucleotides prime synthesis of DNA complementary to at least a portion of said RNA.

10

18. A method as defined in claim 4, wherein in step (a) random hexamer oligonucleotides are also used as reverse transcription primers.

15

20

25

30

35

40

45


50

55

**FIG. 1**

1 - 9


10 - 18




**FIG. 2**

19 - 27

28 - 36



**FIG. 3**



**FIG. 4**

